



## Initial Visit

- Discuss PrEP benefits and possible complications
- Discuss HIV/STI risk reduction
- Screen for acute HIV symptoms. Consider post-exposure prophylaxis (PEP) if high risk exposure in last 72 hours
- Sign PrEP agreement form
- Perform following laboratory tests:
  - HIV test , 4th generation Ag/Ab (+ HIV viral load if concern for acute HIV)
  - STI screening (syphilis & GC/CT NAAT [urine or vagina, rectum, pharynx])
  - Basic metabolic panel
  - HAV total Ab, HBV surface Ag, HBV surface Ab, HBV core Ab, HCV Ab
  - Pregnancy test

After confirmation of negative HIV test, negative pregnancy test, CrCl >60, and negative HBV surface antigen, call and inform patient to start Truvada

**Prescribe 30-day supply, no refills; initiate Truvada as soon as possible, no later than 2 weeks after negative HIV test**

## 30-Day Visit

- Evaluate for eligibility for vaccination to HPV, HAV & HBV
  - HIV test, fourth generation Ag/Ab
  - Evaluate for Truvada side effects
  - Discuss HIV/STI risk reduction
- Prescribe 30-day supply with 2 refills**

### Discuss at Every Visit

- HIV/STI risk reduction
- Provide condoms
- Adherence counseling
- Clean syringe resources
- Evaluate continued PrEP indications

## Every 3-Months

- HIV test
  - Pregnancy test
  - STI screening (syphilis & GC/CT NAAT [urine or vagina, rectum, pharynx])
  - Serum creatinine
  - Discuss HIV/STI risk reduction
- Prescribe 30-day supply with 2 refills**

### Special Considerations for Adolescents

- Monthly visits for at least first 4-6 months
- Provide 30-day supply with no refill during this time
- Consider confidential encounter
- Discuss adherence challenges

## Monitoring Labs

- |                        |  |
|------------------------|--|
| <b>Every 3 months</b>  | <ul style="list-style-type: none"> <li>• HIV test</li> <li>• Assess for STI symptoms</li> <li>• STI screening</li> <li>• Pregnancy test</li> </ul> |
| <b>Every 6 months</b>  | <ul style="list-style-type: none"> <li>• Serum creatinine</li> </ul>   |
| <b>Every 12 months</b> | HCV Ab for MSM, persons who inject drugs, and persons with multiple sex partners   |